

DNA Analysis of Stool for Colon Cancer Screening (Cologuard®)

Policy MP-006

Origination Date: 06/10/2018

Reviewed/Revised Date: 07/17/2024

Next Review Date: 07/17/2025

Current Effective Date: 07/17/2024

Disclaimer:

- 1. Policies are subject to change in accordance with State and Federal notice requirements.
- 2. Policies outline coverage determinations for U of U Health Plans Commercial, CHIP and Healthy U (Medicaid) plans. Refer to the "Policy" section for more information.
- 3. Services requiring prior-authorization may not be covered, if prior-authorization is not obtained.
- 4. This Medical Policy does not guarantee coverage or payment of the service. The service must be a benefit in the member's plan and the member must be eligible for coverage at the time of service. Additional payment guidelines may be applied that are not included in this policy.

Description:

Colorectal cancer (CRC) is one of the most preventable cancers, yet it is second leading cause of cancer death in the United States. In 2021, there will be an estimated 104,270 new cases of colon cancer, 45,230 of rectal cancer and approximately 52,980 people will die from CRC. Without preventive measures, approximately 4-5% of Americans will develop colorectal cancer at some point in their lives. On August 11, 2014, Cologuard[®] was approved by the FDA. This is the first stool-based colorectal screening test that detects the presence of red blood cells and DNA mutations. Cologuard utilizes a multi-target approach to detect DNA and hemoglobin biomarkers associated with colorectal cancer and pre-cancer.

Eleven biomarkers are targeted and provide a stronger connection between colorectal cancer and pre-cancer. Methylation, mutation, and hemoglobin results are combined in the laboratory analysis to provide a single positive or negative reportable result.

Policy Statement and Criteria

1. Commercial Plans/CHIP

U of U Health Plans covers Cologuard[®] for stool colon cancer screening once every 3 years, for average risk members ages 45-75 years when the test is recommended by their physician and it is not within the standard interval of another screening test.

U of U Health Plans does NOT cover any other method of DNA analysis of stool testing for colon cancer screening. Use of this testing is considered investigational/ experimental as current evidence is inadequate to show that stool DNA testing with any test other than Cologuard[®] is an effective way to screen for colon cancer and can improve health outcomes for patients.

2. Medicaid Plans

Coverage is determined by the State of Utah Medicaid program; if Utah State Medicaid has no published coverage position and InterQual criteria are not available, the U of U Health Plans Commercial criteria will apply. For the most up-to-date Medicaid policies and coverage, please visit their website at: <u>https://medicaid.utah.gov/utah-medicaid-official-publications/</u> or the <u>Utah Medicaid code Look-Up tool</u>

CPT/HCPCS codes covered by Utah State Medicaid may still require further evaluation to determine medical necessity for coverage.

Clinical Rationale

As with any diagnostic test, the key outcomes are the diagnostic performance (i.e., sensitivity, specificity, positive and negative predictive value) compared to a gold standard, and consideration of how the results of the test will be used to benefit patient management. Of the various screening options (fecal occult blood testing, flexible sigmoidoscopy, double contrast barium enema, colonoscopy), colonoscopy is considered the gold standard. For example, in patients considered at high risk for colorectal cancer, due either to a family history or hereditary nonpolyposis colorectal cancer (HNPCC) mutation, colonoscopy at varying intervals is recommended by the American Society of Colorectal Surgeons, the American Gastroenterological Society, and the American Cancer Society. Therefore, patients at high risk of colorectal cancer with suspected or known mutations of the HNPCC gene, should have a colonoscopy.

For patients at average risk to moderate risk for colorectal cancer, the above organizations also recommend colonoscopy starting at age 45, with an interval of 10 years, as one screening option. In addition, other screening techniques are also considered options, and the choice of screening option may be dictated in part by patient preference. Many authors have noted the low patient acceptance of current colorectal cancer screening options, particularly flexible sigmoidoscopy and colonoscopy; at the present time only about 40% of eligible patients undergo screening for colon cancer. Advocates of genetic testing of stool samples have hypothesized that the relative simplicity of collecting a stool sample might increase the overall compliance with screening recommendations. Therefore, for patients at average to moderate risk of colon cancer, genetic testing of stool samples will be compared to colonoscopy and also to fecal occult blood testing, the other entirely noninvasive technique. Patient acceptance of the different options is also a relevant outcome as a technique to increase screening compliance.

The available published, peer-reviewed data focus on the technical feasibility of genetic testing of stool samples. For example, Ahlquist and colleagues published a study focusing on the use of a multi-target assay panel for colorectal cancer screening. This retrospective study included 22 patients with known colorectal cancer, 11 with adenomas, and 28 patients with normal colonoscopy examinations. It was not reported whether or not these patients were considered at average, moderate, or high risk for cancer.

The panel included 15 sites on the KRAS gene, p53 and adenomatous polyposis genes, analysis of BAT-26, and highly amplifiable DNA. The panel detected 20 of the 22 cancers (91%) and 9 of the 11 adenomas (82%). The same panel assay was performed on tissue samples from 19 of the 21 cancers. The presence of point mutations was concordant in tissue and stool analysis in 12 of the 19 paired specimens. The authors attributed the high neoplasm detection rate of the stool analysis to the efficient isolation of human DNA from the stool, but also commented that cancers represented in this study were large (median 4 cm in diameter) and symptomatic, and thus may shed more aberrant DNA than smaller cancers. For the 11 patients with adenomas, the results of the stool DNA testing were compared to fecal occult blood testing. While the fecal occult blood testing was negative in all these patients, genetic mutations were detected in the stool sample of all patients with adenomas.

Dong and colleagues performed a study of stool DNA isolated from 51 colorectal cancer patients. The stool DNA and tumor tissue were evaluated for the presence of mutations in the genes p53, BAT-26, and KRAS. The 3 genetic markers together detected 71% of the 51 patients. Of interest, no genetic mutations were identified in the tumor tissue of 15 patients. Other feasibility studies using a variety of markers have also focused on patients with known cancers, and thus these studies do not duplicate the targeted populations for screening. No prospective studies were found in the published literature comparing the diagnostic performance of analysis of DNA from stool samples to either colonoscopy or fecal occult blood testing among either average to moderate risk to high risk patients. For average risk patients, the published feasibility studies focused on the use of different panels of DNA markers. No study identified focused on the use of the single marker, BAT-26, in patients with known or suspected mutations of the HNPCC gene. No studies discussed how the use of DNA analysis in stool samples might supplant or enhance current screening options.

Notably, the study by Imperiale et al., was a non-randomized, cross-sectional, multicenter trial of 9,989 patients who were included in the primary analysis. Results from a single, albeit large, study show that Cologuard has better sensitivity and worse specificity than FIT across various clinical manifestations. The other 2 included studies are 1) a model that is based on incorrect pricing information and 2) a study unique to a specific population not relevant in a broad sense. Though the test has a recommended use of once every 3 years, 1 systematic review illustrated its benefit is best if used annually. This may impact its cost effectiveness in real world settings.

Hayes updated their tech assessment for the use of Cologuard when screening for colorectal cancer (CRC) in average risk adults on March 24, 2023. They found no studies for analytical validity, 11 studies for clinical validity, and 3 studies for clinical utility. The evidence for clinical validity suggests that Cologuard's ability to accurately detect CRC may be comparable to that of fecal immunochemical testing (FIT). However, the ability of Cologuard to detect a truly negative test is unclear. Few studies provided statistical comparisons with FIT; although, numerically, Cologuard appears to provide more false positives than FIT. A single study comparing the accuracy of Cologuard with virtual computed tomography (CT) colonography found CT colonography accurately identified more CRC, advanced lesions and high-risk sessile serrated polyps (SSPs), suggesting that CT outperformed Cologuard. When repeating Cologuard 3 years after the initial test, the predictive power to detect advanced lesions remained the same. Comparing Cologuard with colonoscopy alone, following a positive Cologuard test, Cologuard identified statistically more neoplasia than colonoscopy alone (Cologuard, 67%; colonoscopy alone, 41%; P<0.0001). However, evidence of clinical utility is insufficient to draw conclusions on the effect of the Cologuard test on follow-up colonoscopy and patient outcomes. The assessment concluded that the primary support for the Cologuard test comes from evidence suggesting that Cologuard is comparable to FIT in its ability to detect CRC and advanced precancerous lesions. The evidence supporting Cologuard's ability to detect a truly negative test compared with FIT remains unclear due to

inconsistency in results and a lack of statistical comparisons. As for clinical utility there is insufficient evidence due to poor quality of studies and lack of consistency.

A single, small study evaluating change in patient management and 2 test preference studies do not provide sufficient evidence of clinical utility. Furthermore, these studies do not demonstrate improved outcomes. In the future, blood-based CRC screening may have a role in CRC screening, such as for patients who decline the usual recommended testing techniques; however, there is currently not enough evidence demonstrating a positive change in management or an improvement in outcomes to justify its use without additional studies.

In 2024, UpToDate reviewed updated and current literature for the screening of colorectal cancer by multi-targeted stool DNA (mt-sDNA) tests with fecal immunochemical testing (Cologuard). The evidence of effectiveness of mt-sDNA is based on comparison with studies of other screening strategies and modelling studies. No randomized trials of mt-sDNA for screening for CRC were found. Sensitivity for CRC appears to exceed that for FIT, no matter what stage. However, evidence for improved clinical outcomes is indirect. Also noted is that stool-based screening tests for colon cancer have a higher detection rate for CRC and less so in detecting precancerous adenoma polyps. Every positive CRC screening test other than colonoscopy should be followed immediately by a colonoscopy. A negative screening result should be re-tested for the recommended interval of every 3 years.

The American College of Gastroenterology (ACG) is currently reviewing their existing colorectal cancer screening guidelines, the current recommendations are as follows:

- Tests that prevent cancer are preferred over those that only detect cancer;
- The preferred colorectal cancer prevention test is colonoscopy every 10 years beginning at age 45;
- For patients who decline colonoscopy or another cancer prevention test, the preferred cancer test is FIT to be conducted annually (alternative cancer detection tests include annual Hemoccult Sensa® and fecal DNA testing every 3 years).

The National Comprehensive Cancer Network (NCCN) CRC screening guidelines (Version 1.2024) recommend the inclusion of mt-sDNA-based testing as a potential screening modality in average risk patients aged ≥45. However, data to help determine screening intervals and how mt-sDNA testing may fit into an overall screening program are limited. Intervals of every 3 years has been recommended by the FDA. In addition, there are no or limited data in high-risk individuals who refuse colonoscopy or have limited access to conventional screening strategies; therefore, the use of mt-sDNA–based testing should be individualized in these cases.

Applicable Coding

CPT Codes

81528 Oncology (colorectal) screening, quantitative real-time target and signal amplification of 10 DNA markers (KRAS mutations, promoter methylation of NDRG4 and BMP3) and fecal hemoglobin, utilizing stool, algorithm reported as a positive or negative result (Cologuard[®])

HCPCS Codes

No applicable codes

References:

- 1. Ahlquist, D. A. (2015). Multi-target stool DNA test: a new high bar for noninvasive screening. Digestive diseases and sciences, 60(3), 623-633.
- 2. Ahlquist DA, Skoletsky JE, Boynton KA et al. (2000). Colorectal cancer screening by detection of altered human DNA in stool: feasibility of a multitarget assay panel. Gastroenterology;119(5):1219-27
- 3. American Cancer Society. Colorectal Cancer. (2021) "Key Statistics for Colorectal Cancer" Last revised: January 12, 2021. Accessed: May 25, 2021. Available at: <u>https://www.cancer.org/cancer/colon-rectal-cancer/about/key-statistics.html</u>
- 4. Berger, B.M., USPSTF Colorectal Cancer Screening Guidelines: An Extended Look at Multi-Year Interval Testing. American Journal of Managed Care, 2016. 22(2): p. 77-81.
- 5. Centers for Disease Control and Prevention (CDC). Colorectal (Colon) Cancer. Last reviewed: May 24, 2021. Accessed May 25, 2021. Available at: https://www.cdc.gov/cancer/colorectal/
- 6. Dong SM, Traverso G, Johnson C et al. (2001). Detecting colorectal cancer in stool with the use of multiple genetic targets. J Natl Cancer Inst;93(11):858-65.
- Doubeni, C., et al. (2022) UpToDate "Tests for screening for colorectal cancer" Topic: 7576. Version 110.0. Last updated May 2024. Literature review current through April 2024. Accessed: May 20, 2024. Available at: <u>https://www.uptodate.com/contents/tests-for-screening-for-colorectalcancer?topicRef=2496&source=see_link#H240481080</u>
- 8. Hayes, Inc. (2021) Molecular Test Assessment "Cologuard (Exact Science Corp.)" Last Reviewed: March 13, 2024. Accessed: May 20, 2024. Available at: <u>https://evidence.hayesinc.com/report/gte.cologuard3111</u>
- 9. Imperiale, T.F., et al., Multitarget Stool DNA Testing for Colorectal-Cancer Screening. New England Journal of Medicine, 2014. 370(14): p. 1287-1297.
- 10. Koshiji M, Yonekura Y, Siato T et al. (2002). Microsatellite analysis of fecal DNA for colorectal cancer detection. J Surg Oncol;80(1):34-40
- 11. Levin B, Brooks D, Smith RA et al. (2003). Emerging technologies in screening for colorectal cancer: CT colonography, immunochemical fecal occult blood tests, and stool screening using molecular markers. CA Cancer J Clin; 53(1):44-55
- 12. Levin B, Lieberman DA, McFarland B, et al. (2008). Screening and surveillance for the early detection of colorectal cancer and adenomatous polyps, 2008: A joint guideline from the American Cancer Society, the US Multi-Society Task Force on colorectal cancer, and the American College of Radiology. CA Cancer Journal for Clinicians, 58 (3), 130-160
- 13. National Comprehensive Cancer Network (NCCN) "Guidelines Version 1.2024 Colorectal Cancer Screening" February 27, 2024. Accessed May 20, 2024. Available at: <u>https://www.nccn.org/professionals/physician_gls/pdf/colorectal_screening.pdf</u>
- 14. Redwood, D.G., et al., Stool DNA Testing for Screening Detection of Colorectal Neoplasia in Alaska Native People. Mayo Clin Proc, 2016. 91(1): p. 61-70.
- 15. The American Journal of Gastroenterology (AJG) (2008) "American College of Gastroenterology Guidelines for Colorectal Cancer Screening" Accessed: May 25, 2021. Available at:

https://journals.lww.com/ajg/Fulltext/2009/03000/American_College_of_Gastroenterology_Guidelines.36.aspx

- 16. Traverso G, Shuber A, Olsson L et al. (2002). Detection of proximal colorectal cancers through analysis of fecal DNA. Lancet;359(9304):403-4
- 17. U.S Preventive Services Task Force (USPFTF). Colorectal Cancer: Screening. May 18, 2021. Accessed May 20, 2021. Available at: https://www.uspreventiveservicestaskforce.org/uspstf/recommendation/colorectal-cancer-screening
- 18. Winawer S, Fletcher R, Rex D, et al. (2003). Colorectal cancer screening and surveillance: clinical guidelines and rationale-Update based on new evidence. Gastroenterology; 124(2):544-60.

Disclaimer:

This document is for informational purposes only and should not be relied on in the diagnosis and care of individual patients. Medical and Coding/Reimbursement policies do not constitute medical advice, plan preauthorization, certification, an explanation of benefits, or a contract. Members should consult with appropriate health care providers to obtain needed medical advice, care, and treatment. Benefits and eligibility are determined before medical guidelines and payment guidelines are applied. Benefits are determined by the member's individual benefit plan that is in effect at the time services are rendered.

The codes for treatments and procedures applicable to this policy are included for informational purposes. Inclusion or exclusion of a procedure, diagnosis or device code(s) does not constitute or imply member coverage or provider reimbursement policy. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage of these services as it applies to an individual member.

U of U Health Plans makes no representations and accepts no liability with respect to the content of any external information cited or relied upon in this policy. U of U Health Plans updates its Coverage Policies regularly, and reserves the right to amend these policies and give notice in accordance with State and Federal requirements.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from U of U Health Plans.

"University of Utah Health Plans" and its accompanying logo, and its accompanying marks are protected and registered trademarks of the provider of this Service and or University of Utah Health. Also, the content of this Service is proprietary and is protected by copyright. You may access the copyrighted content of this Service only for purposes set forth in these Conditions of Use.

© CPT Only – American Medical Association