

Implantable Hemodynamic Monitor for Managing Patients with Heart Failure (e.g. CardioMEMS)

Policy MP-035

Origination Date: 01/30/2019

Reviewed/Revised Date: 12/18/2025

Next Review Date: 12/18/2026

Current Effective Date: 12/18/2025

Disclaimer:

1. Policies are subject to change in accordance with State and Federal notice requirements.
2. Policies outline coverage determinations for U of U Health Plans Commercial, CHIP, Healthy U (Medicaid) and Health Choice Utah (Medicaid) plans. Refer to the "Policy" section for more information.
3. Services requiring prior-authorization may not be covered, if prior-authorization is not obtained.
4. **This Medical Policy does not guarantee coverage or payment of the service. The service must be a benefit in the member's plan, and the member must be eligible for coverage at the time of service. Additional payment guidelines may be applied that are not included in this policy.**

Description:

According to the American Heart Association (AHA), heart failure (HF) is a chronic, progressive condition in which the body does not receive enough oxygen because the heart muscle is unable to pump enough blood throughout the body to meet its oxygen needs. Signs and symptoms of HF, as described by the National Heart, Lung, and Blood Institute (NHLBI), include trouble breathing or shortness of breath, fatigue, and swelling in the hands, feet, abdomen, legs, and neck veins. These symptoms tend to worsen as the disease progresses. As blood flow to the kidneys decreases, they begin to retain sodium, which causes the body to accumulate fluid. When the body begins to build up excess fluid, other symptoms such as weight gain, frequent urination, and coughing may also develop due to fluid buildup in the lungs. If this is associated with HF, it could indicate pulmonary edema, which requires emergency treatment.

The New York Heart Association (NYHA) categorizes heart failure (HF) using a functional classification system based on limitations in physical activity limits and the symptomatic status of HF (*see* Table A). The American College of Cardiology Foundation (ACCF) and the American Heart Association (AHA) classify HF based on its development and progression (*see* Table B).

Approximately 5.7 million people in America will be affected with HF and about half of those will die around 5 years after they are diagnosed. There is no cure, it affects children and adults, generally adults 65 and older. The cost to treat HF is estimated at \$30 billion a year when you

include all health-related expenses including: lost workdays, healthcare services, and pharmacological treatment associated with HF management (CDC, 2016).

Standard management of HF is to regularly monitor the signs and symptoms such as weight gain, swelling, fatigue, and shortness of breath. CardioMEMS (Abbott [acquired from St. Jude Medical Inc. in 2017]) is a minimally invasive, wireless implantable hemodynamic monitoring device that provides continuous monitoring of pulmonary artery pressure (PAP) for use at home in patients with HF. This device offers an aid for early detection and prevention of HF decompensation with a PAP sensor, a transvenous catheter delivery system, a patient home-monitoring electronic system, and a secure internet accessible database that allows clinicians to access patient data.

Policy Statement and Criteria

1. Commercial Plans/CHIP

U of U Health Plans does NOT cover implantable hemodynamic monitors (e.g. CardioMEMS) for the management of patients with NYHA * Class III heart failure as the safety and efficacy of this device has not been established.

***Table A - NYHA Heart Failure Functional Classification**

NYHA Class	Limitations on Physical Activity	Impact on Physical Activity
I	No limitations	Ordinary physical activity does not cause undue fatigue, palpitation, or dyspnea
II	Slight limitation	Comfortable at rest. Ordinary physical activity results in fatigue, palpitation, or dyspnea.
III	Marked limitation	Comfortable at rest. Less than ordinary activity causes fatigue, palpitation, or dyspnea.
IV	Unable to carry on any physical activity without discomfort	Symptoms of heart failure at rest. If any physical activity is undertaken, discomfort increases.

Table B – ACCF and AHA Heart Failure Class System

Class	Symptoms
Class A	No objective evidence of cardiovascular disease. No symptoms and no limitation in ordinary physical activity.
Class B	Objective evidence of minimal cardiovascular disease. Mild symptoms and slight limitation during ordinary activity. Comfortable at rest.
Class C	Objective evidence of moderately severe cardiovascular disease. Marked limitation in activity due to symptoms, even during less-than-ordinary activity. Comfortable only at rest.
Class D	Objective evidence of severe cardiovascular disease. Severe limitations. Experiences symptoms even while at rest.

2. Medicaid Plans

Coverage is determined by the State of Utah Medicaid program; if Utah State Medicaid has no published coverage position and InterQual criteria are not available, the U of U Health Plans Commercial criteria will apply. For the most up-to-date Medicaid policies

and coverage, please visit their website at: <https://medicaid.utah.gov/utah-medicaid-official-publications/> or the [Utah Medicaid code Look-Up tool](#)

CPT/HCPCS codes covered by Utah State Medicaid may still require further evaluation to determine medical necessity for coverage.

Clinical Rationale

The CHAMPION trial, a 2011 single-blind trial (Abraham et al) found that the CardioMEMS implantable hemodynamic monitoring system reduced the rate of hospital admissions. Sixty-four centers in the U.S were enrolled with patients that had a previous hospital admission for heart failure (HF) and were diagnosed with NYHA class III heart failure, irrespective of the left ventricular ejection fraction. Patients were randomly assigned to their group and masked as to which group they belonged. The treatment group was managed with a wireless implantable hemodynamic monitoring (W-IHM) system and the clinicians used daily measurement of pulmonary artery pressures in addition to standard of care. The control group experienced standard of care alone, both groups were treated for at least 6 months. The primary efficacy endpoint was the rate of heart-failure-related hospitalizations at 6 months. The safety endpoints assessed at 6 months were freedom from device-related or system-related complications (DSRC) and freedom from pressure-sensor failures. All analyses were by intention to treat. In 6 months, 83 heart-failure-related hospitalizations were reported in the treatment group (n = 270) compared with 120 in the control group (n = 280; rate 0.31 versus 0.44, hazard ratio [HR] 0.70, 95% CI: 0.60 to 0.84, p < 0.0001). During the entire follow-up (mean 15 months [SD 7]), the treatment group had a 39% reduction in heart-failure-related hospitalization compared with the control group (153 versus 253, HR 0.64, 95% CI: 0.55 to 0.75; p < 0.0001). Eight patients had DSRC and overall freedom from DSRC was 98.6% (97.3 to 99.4) compared with a pre-specified performance criterion of 80% (p < 0.0001); and overall freedom from pressure-sensor failures was 100% (99.3 to 100.0).

A 2015 retrospective analysis (Benza et al) of the CHAMPION trial examined the implantable hemodynamic monitor (IHM) in 550 NYHA Functional Class III HF patients, regardless of left ventricular ejection fraction (LVEF) or HF etiology. Evaluations of clinical variables, changes in medical therapy, HF hospitalization rates and survival in patients with and without World Health Organization (WHO) Group II pulmonary hypertension (PH) were conducted. Data were obtained for 314 patients (59%) who had WHO Group II PH. Patients without PH were at significantly lower risk for mortality than PH patients (hazard ratio [HR] 0.31, 95% CI: 0.19 to 0.52, p < 0.0001). Pulmonary hypertension patients had higher HF hospitalization rates than non-PH patients (0.77/year versus 0.37/year; HR 0.49, 95% CI: 0.39 to 0.61, p < 0.001). In patients with and without PH, ongoing knowledge of hemodynamic data resulted in a reduction in HF hospitalization for PH patients (HR 0.64, 95% CI: 0.51 to 0.81, p = 0.002) and for non-PH patients (HR 0.60, 95% CI: 0.41 to 0.89, p = 0.01). Among PH patients, there was a reduction in the composite endpoint of death and HF hospitalization with ongoing knowledge of hemodynamics (HR 0.74, 95% CI: 0.55 to 0.99, p = 0.04), but no difference in survival (HR 0.78, 95% CI: 0.50 to 1.22, p = 0.28). In conclusion WHO Group II PH associated with left heart disease is quite prevalent and identifies HF patients at risk for adverse outcomes. Ongoing knowledge of hemodynamic variables at frequent time intervals from the patient's home, which cannot practically be provided by right heart catheterization (RHC), may allow for more effective treatment strategies to reduce the morbidity of this disease.

Within the CHAMPION trial, a 2016 subgroup analysis by Costanzo et al, compared medical therapy in patients who were managed with CardioMEMS implantable hemodynamic monitors (CM-IHM) and those who received standard management only. In patients managed with CM-IHM for HF (Heart

Failure)-related hospitalizations, there was a reduction of stay, the subgroup did an analysis to determine what medical interventions appeared to produce these reductions. Baseline medical therapy was similar among patient groups; however, patients managed with CM-IHM had a significantly higher frequency of medication adjustments, particularly in the number of diuretic (1547 for CM-IHM versus 585 for SM; $P < 0.001$) and vasodilator (293 for CM-IHM versus 104 for SM; $P < 0.05$) dose changes. These findings suggest an inverse relationship between increased medication management and HF rates among patients whose treatment was managed by CM-IHM data.

In a 2017 retrospective cohort study (Desai et al) of 1935 Medicare enrollees who underwent implantation of the CM-IHM following FDA approval, 1114 individuals were continuously enrolled and had evaluable Medicare administrative claims data for at least 6 months before and following implantation to use for the study. A subset of 480 enrollees had complete data for 12 months before and after implantation. The cumulative incidence of HF-related hospitalizations were significantly lower in the post-implantation period than in the preimplantation period at both 6- and 12-month follow-ups. Limitations of this pre-post retrospective study include lack of data on medical history, ejection fraction, indication for implantation and possible confounding due to amplified touchpoints with the health care system necessitated by the device's implantation.

In 2018, Kilic and colleagues examined if there is definitive criteria to establish patient selection for monitoring HF in patients with a CM-IHM. Clinical trials such as the CHAMPION RCT suggest that patients with moderate HF on optimal medical therapy with difficult-to-control volume status and frequent hospitalizations for decompensated HF appear to achieve the most benefit from pressure-guided HF monitoring. Factors such as ejection fraction, PH, or COPD do not seem to suggest an impact of the benefit from CM-IHM on reducing HF-related hospitalization according to a subgroup analyses using data from the CHAMPION RCT. Findings in patients with HF who were not part of a clinical trial, who were included in a large registry analysis, suggested that the benefits of CM-IHM may extend to the general population of patients with HF. Results of another registry analysis ($n=430$ pts) demonstrated that patients monitored with CM-IHM who subsequently had a LVAD implanted had higher mean PAP levels in the months leading up to LVAD placement compared with patients who did not receive LVAD placement, suggesting that CM-IHM may aid in clinical decision making about the timing of LVAD implant and the management of LVAD patients. In conclusion, the authors found that further studies are needed to establish definitive patients' selection criteria for the monitoring HF with CM-IHM, as currently there is not enough data to determine this criteria.

On March 3, 2023, St. Jude Medical initiated a recall which remains open as of December 13, 2024. They noted that "Radiated Radiofrequency (RF) emissions of the CardioMEMS Patient Electronics Systems (PES) Models CM1000, CM1010 and CM1100 and CardioMEMS Hospital Electronics Systems (HES) Model CM3000 exceed certain emissions limits and have the potential to cause interference with other medical devices, such as implantable neurostimulators, pacemakers, cardiac defibrillators, continuous glucose monitors, or other bio-wearable sensors when in close proximity to an active Hospital or Patient Electronics System (i.e., during readings)." (Recall number Z-1219-2023).

The CardioMems is a percutaneous sensor placed into the pulmonary artery and is utilized for continuous hemodynamic monitoring of those being managed for CHF (Yancy et al, 2017). It has been developed as a means to assist in the treatment of such individuals. Evaluation of this device has consisted of single-blinded trials, as well as some retrospective and sub- group analyses. These assessments have essentially demonstrated that the device may decrease hospitalizations for those with congestive heart failure. Specifically, those with more severe CHF (NYHA class III) and/or those who have been hospitalized more frequently experienced a lower hospitalization rate. A sub-group analysis of the CHAMPION Trial demonstrated that those with the device had a higher frequency of medication

adjustments, which possibly lead to a lower hospitalization rate. However, as these were basically retrospective, single-blind, and sub- group analyses, a benefit of the device towards overall improved cardiac outcomes has not been definitively established. Additionally, a mortality benefit or long-term quality of life benefit has also not been established. Thus, there is no definitive evidence that the monitor, compared to conventional medical management and clinical monitoring, has been established as standard of care towards improved overall CHF or cardiac outcomes (morbidity and mortality). As such the plan criteria appropriately does not cover the device for the management of patients with NYHA class III heart failure as the safety and efficacy of this device has not been established.

Hayes updated their health technology assessment on the CardioMEMS implantable hemodynamic monitor (CM-IHM) for managing patients with heart failure in July of 2022. There continues to be a very-low-quality body of evidence that is insufficient to draw conclusions regarding the safety and efficacy of the CM-IHM in adults with NYHA class III HF. Limited evidence suggested management of HF patients with CM-IHM in addition to standard care, reduced the incidence of HF-related hospitalization. In conclusion, the authors found that due to inconsistent findings, short duration of follow-up, and variability in reported outcome measures, substantial uncertainty remains about the comparative effectiveness of CM-IHM with standard monitoring and the impact of CM-IHM on long-term patient's safety, mortality, survival, and quality of life. Thus, due to the uncertainty surrounding the efficacy of CM-IHM, large well-designed comparative studies are needed to provide evidence for the long-term benefits and harms of CM-IHM.

Hayes updated their research brief regarding CM-IHM for Managing Patients with Heart Failure in December of 2021 to include 7 newly published studies (1 RCT and 6 single-arm studies). Interventions included CM-IHM, CM-IHM plus HeartLogic and CM-IHM plus sacubitril/valsartan. The RCT compared CM-IHM with control. Other comparisons included sex, body mass index (BMI), ejection fraction, and pulmonary artery proportional pulse pressure (PAPP). Outcomes evaluated the effect on comorbidities, cardiac hospitalizations, use of diuretics, and mortality. Of note, nearly all of the trials were either funded by Abbott or included physicians with possible conflict of interests related to their affiliation with Abbott. The authors concluded that further unbiased research is still warranted to validate these findings and demonstrate clinical significance.

Applicable Coding

CPT Codes

93701 Bioimpedance-derived physiologic cardiovascular analysis

93799 Unlisted cardiovascular service or procedure

HCPCS Codes

No applicable codes

References:

1. Abbott. CardioMEMS™ HF System Program Practice Guide. 2017. Available at: <https://www.sjm.com/en/professionals/education-training/professional-education/hf-management/cardiomems-workflow-resources>. Accessed January 12, 2019.
2. Abbott. St. Jude Medical CardioMEMS HF System is Now Guideline Directed Therapy for Heart Failure Patients in Europe [news release]. June 6, 2016. Available at: <http://media.sjm.com/newsroom/news-releases/news-releases-details/2016/St-Jude-Medical-CardioMEMS-HF-System-is-Now-Guideline-Directed-Therapy-for-Heart-Failure-Patients-in-Europe/default.aspx>. Accessed January 12, 2019.
3. Abraham WT, Adamson PB, Bourge RC, et al.; CHAMPION Trial Study Group. Wireless pulmonary artery haemodynamic monitoring in chronic heart failure: A randomised controlled trial. *Lancet*. 2011;377(9766):658-666.

4. American Heart Association (AHA) Health Topics-Heart Failure-"What is Heart Failure?" Last reviewed May 31, 2017. Assessed Jan 11, 2019. Available at: <https://www.heart.org/en/health-topics/heart-failure/what-is-heart-failure>
5. American Heart Association (AHA) Health Topics-Heart Failure-What is Heart Failure?-"Classes of Heart Failure" (NYHA Heart Failure Functional Classification and ACCF and AHA Heart Failure Class System). Last reviewed: May 31, 2017. Assessed: Jan 16, 2019. Available at: <https://www.heart.org/en/health-topics/heart-failure/what-is-heart-failure/classes-of-heart-failure>
6. American Heart Association (AHA). Pulmonary Hypertension – High Blood Pressure in the Heart-to-Lung System. Updated October 31, 2016. Available at: http://www.heart.org/HEARTORG/Conditions/HighBloodPressure/AboutHighBloodPressure/What-is-Pulmonary-Hypertension_UCM_301792_Article.jsp. Accessed January 14, 2019.
7. American Heart Association (AHA). Types of Heart Failure. May 31, 2017a. Available at: <https://www.heart.org/en/health-topics/heart-failure/what-is-heart-failure/types-of-heart-failure>. Accessed January 14, 2019.
8. American Heart Association (AHA). Your Heart Failure Healthcare Team. Updated May 31, 2017b. Available at: <http://www.heart.org/en/health-topics/heart-failure/living-with-heart-failure-and-managing-advanced-hf/your-heart-failure-healthcare-team>. Accessed January 14, 2019.
9. Ayyadurai P, Alkhwam H, Saad M, et al. An update on the CardioMEMS pulmonary artery pressure sensor. *Ther Adv Cardiovasc Dis*. 2019;13:1753944719826826. doi:10.1177/1753944719826826
10. Benza RL, Raina A, Abraham WT, et al. Pulmonary hypertension related to left heart disease: Insight from a wireless implantable hemodynamic monitor. *J Heart Lung Transplant*. 2015;34(3):329-337.
11. Brugts JJ, Radhoe SP, Clephas PR, et al. Remote haemodynamic monitoring of pulmonary artery pressures in patients with chronic heart failure (MONITOR-HF): a randomised clinical trial. *Lancet*. 2023;401(10394):2113-2123. doi:10.1016/s0140-6736(23)00923-6
12. Centers for Disease Control and Prevention (CDC). Heart Failure Fact Sheet. Updated June 16, 2016. Available at: http://www.cdc.gov/dhdsp/data_statistics/fact_sheets/fs_heart_failure.htm. Accessed January 14, 2019.
13. Clyde W. Yancy , Mariell Jessup , Biykem Bozkurt , Javed Butler , Donald E. Casey , Monica M. Colvin , Mark H. Drazner , Gerasimos S. Filippatos , Gregg C. Fonarow , Michael M. Givertz , Steven M. Hollenberg , JoAnn Lindenfeld , Frederick A. Masoudi , Patrick E. McBride , Pamela N. Peterson , Lynne Warner Stevenson , and Cheryl Westlake. (2017) "ACC/AHA/HFSA Focused Update of the 2013 ACCF/AHA Guideline for the Management of Heart Failure: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Failure Society of America". Originally published 28 Apr 2017 *Circulation*. 2017;136:e137–e161.
14. Costanzo MR, Stevenson LW, Adamson PB, et al. Interventions linked to decreased heart failure hospitalizations during ambulatory pulmonary artery pressure monitoring. *JACC Heart Fail*. 2016;4(5):333-344.
15. Cowie MR, Flett A, Cowburn P, Foley P, Chandrasekaran B, Loke I, Critoph C, Gardner RS, Guha K, Betts TR, Carr-White G. Real-world evidence in a national health service: results of the UK CardioMEMS HF system Post-Market study. *ESC heart failure*. 2022 Feb;9(1):48-56.
16. Desai AS, Bhimaraj A, Bharmi R, et al. Ambulatory Hemodynamic Monitoring Reduces Heart Failure Hospitalizations in "Real-World" Clinical Practice. *J Am Coll Cardiol*. May 16, 2017;69(19):2357-2365. PMID 28330751
17. Food and Drug Administration (FDA). CardioMEMSTM HF System Patient Electronics System (PES) – P100045. March 3, 2023. Accessed: December 13, 2024. Available at: <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfres/res.cfm?id=198347>.
18. Hayes, Winifred S., Inc. (2019). A Health Technology Brief/Evidence Analysis Research Brief "CardioMEMS Implantable Hemodynamic Monitor (Abbott) for Managing Patients with Heart Failure" Updated: July 28, 2022. Last review: August 1, 2024. Assessed: December 10, 2024. Available at: https://evidence.hayesinc.com/report/htb_wireless3228
19. Kilic A, Katz JN, Joseph SM, et al.; Evolving Mechanical Support Research Group (EMERG) Investigators. Changes in pulmonary artery pressure before and after left ventricular assist device implantation in patients utilizing remote haemodynamic monitoring. *ESC Heart Fail*. 2018. Epub ahead of print October 23, 2018. Available at: <https://doi.org/10.1002/ehf2.12373>. Accessed January 15, 2019.
20. National Heart, Lung, and Blood Institute (NHLBI). Signs, Symptoms, and Complications [of HF]. 2018b. Available at: <https://www.nhlbi.nih.gov/health-topics/heart-failure>. Accessed January 14, 2019.
21. Cowie MR, Flett A, Cowburn P, Foley P, Chandrasekaran B, Loke I, Critoph C, Gardner RS, Guha K, Betts TR, Carr-White G. Real-world evidence in a national health service: results of the UK CardioMEMS HF system Post-Market study. *ESC heart failure*. 2022 Feb;9(1):48-56.
22. Yancy, C. W., et al. (2017). "2017 ACC/AHA/HFSA Focused Update of the 2013 ACCF/AHA Guideline for the Management of Heart Failure: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Failure Society of America." *Circulation* 136(6): e137-e161.

Disclaimer:

This document is for informational purposes only and should not be relied on in the diagnosis and care of individual patients. Medical and Coding/Reimbursement policies do not constitute medical advice, plan preauthorization, certification, an

explanation of benefits, or a contract. Members should consult with appropriate health care providers to obtain needed medical advice, care, and treatment. Benefits and eligibility are determined before medical guidelines and payment guidelines are applied. Benefits are determined by the members' individual benefit plan that is in effect at the time services are rendered.

The codes for treatments and procedures applicable to this policy are included for informational purposes. Inclusion or exclusion of a procedure, diagnosis or device code(s) does not constitute or imply member coverage or provider reimbursement policy. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage of these services as it applies to an individual member.

U of U Health Plans makes no representations and accepts no liability with respect to the content of any external information cited or relied upon in this policy. U of U Health Plans updates its Coverage Policies regularly and reserves the right to amend these policies and give notice in accordance with State and Federal requirements.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from U of U Health Plans.

"University of Utah Health Plans" and its accompanying logo, and its accompanying marks are protected and registered trademarks of the provider of this Service and or University of Utah Health. Also, the content of this Service is proprietary and is protected by copyright. You may access the copyrighted content of this Service only for purposes set forth in these Conditions of Use.

© CPT Only – American Medical Association