

Fecal Microbiota Transplant

Policy MP-062

Origination Date: 09/10/2020

Reviewed/Revised Date: 07/17/2024

Next Review Date: 07/17/2025

Current Effective Date: 07/17/2024

Disclaimer:

1. Policies are subject to change in accordance with State and Federal notice requirements.
2. Policies outline coverage determinations for U of U Health Plans Commercial, CHIP and Healthy U (Medicaid) plans. Refer to the "Policy" section for more information.
3. Services requiring prior-authorization may not be covered, if prior-authorization is not obtained.
4. **This Medical Policy does not guarantee coverage or payment of the service. The service must be a benefit in the member's plan and the member must be eligible for coverage at the time of service. Additional payment guidelines may be applied that are not included in this policy.**

Description:

Clostridium difficile infection (CDI), is a Gram-positive, spore-forming bacterium usually spread by the fecal-oral route. Patients with recurrent CDI have been observed to have reduced diversity of the intestinal microbiome and diminished numbers of bacteria relative to healthy individuals that can cause disease asymptomatic carriage, mild diarrhea, colitis, or pseudomembranous colitis.

According to the American Journal of Gastroenterology (AJG) Clostridium difficile infection is a leading cause of hospital-associated gastrointestinal illness. Patients typically have extended lengths-of-stay and CDI is a frequent cause of large hospital outbreaks of disease.

Fecal microbiota transplant (FMT) is the term used when stool is taken from a healthy individual and instilled into a sick person with certain conditions, such as recurrent CDI. Studies show that patients with recalcitrant CDI have abnormally proportioned colon microbiota and that reintroduction of normal bacteria via donor feces corrects this imbalance and breaks the cycle of CDI recurrence. The purpose of FMT treatment is based on the premise that an imbalance in the community of microorganisms residing in the gastrointestinal (GI) tract is associated with specific disease states, including susceptibility to infection. In its healthy state, intestinal microbiota performs a variety of useful functions including aiding in the digestion of carbohydrates, repressing the growth of pathogenic microbes, mediating the synthesis of certain vitamins, and stimulating the lymphoid tissue to produce antibodies to pathogens.

FMT may be administered by oral capsules, colonoscopy, retention enema, or through a nasojejunal (NJ)/nasoduodenal (ND) tube in the upper GI tract. The choice of the delivery route

depends in part on patient preferences, individual risk, cost, availability of resources and expertise.

Policy Statement and Criteria

1. Commercial Plans/CHIP

U of U Health Plans considers fecal microbiota transplantation including capsulized, frozen and suspension (via rectal enema), medically necessary for treatment of patients with recurrent *Clostridium difficile* infection when the following criteria are met:

- A. At least 3 episodes of recurrent mild to moderate CDI* and failure of a 6-8 week taper with vancomycin with or without an alternative antibiotic (e.g., rifaximin, nitazoxanide, metronidazole, vancomycin); **and**
- B. Persistent positive stool *C. difficile* toxin by testing and at least **one** of the following:
 - i. At least two episodes of recurrent severe CDI[#] resulting in hospitalization and associated significant morbidity, including renal failure.
 - ii. Moderate CDI not responding to standard therapy (vancomycin) for at least a 10 days.
 - iii. Severe fulminant *C. difficile* colitis with no response to standard therapy after 48 hours.

*Patients with moderate clinical disease have frequent loose, bloody stools (>4 per day), mild anemia not requiring blood transfusions, and abdominal pain that is not severe. Patients have minimal signs of systemic toxicity, including a low-grade fever. Adequate nutrition is usually maintained, and weight loss is not associated with moderate clinical disease.

[#]Patients with a severe clinical presentation typically have frequent loose, bloody stools (≥6 per day) with severe cramps and evidence of systemic toxicity as demonstrated by a fever (temperature ≥37.5°C), tachycardia (HR ≥90 beats/minute), anemia (hemoglobin <10.5 g/dL), or an elevated ESR (≥30 mm/hour). Patients may have rapid weight loss.

U of U Health Plans considers fecal microbiota transplantation investigational/experimental in all other circumstances.

U of U Health Plans does NOT cover Rebyota™ suspension as current evidence is insufficient to show superiority to standard fecal microbiota preparations.

2. Medicaid Plans

Coverage is determined by the State of Utah Medicaid program; if Utah State Medicaid has no published coverage position and InterQual criteria are not available, the U of U Health Plans Commercial criteria will apply. For the most up-to-date Medicaid policies

and coverage, please visit their website at: <https://medicaid.utah.gov/utah-medicaid-official-publications/> or the [Utah Medicaid code Look-Up tool](#)

CPT/HCPCS codes covered by Utah State Medicaid may still require further evaluation to determine medical necessity for coverage.

Clinical Rationale

A 2017 systematic review (Quraishi et al) investigated the effect of FMT in patients with recurrent or refractory CDI. In the pooled analysis, 92% of patients had a resolution of CDI (95% CI, 89% to 94%); heterogeneity was classified as likely moderate ($I^2=59\%$). Additionally, in the 7 trials that evaluated FMT, the intervention overall was associated with an increase in the resolution of recurrent and refractory CDI (relative risk, 0.23; 95% CI, 0.07 to 0.80). The 30 case series reported resolution rates for CDI ranged from 68% to 100%. The reviewers considered the RCTs as having a low risk of bias for adequate randomization with allocation concealment and intention-to-treat analysis. Nor did they report an assessment of bias in terms of blinding, sample size adequacy, or possible differences in baseline characteristics. However, they disputed that none of the trials demonstrated the efficacy of FMT as being truly placebo-controlled, and the case series followed patients until resolution of CDI (10 weeks to 8 years), though some had an insufficient follow-up.

In another 2018 meta-analysis (Khan et al.) researched the literature of pooled data on the use of FMT as a treatment option for recurrent CDI. Reviewers only selected randomized controlled trials that compared FMT (fresh or frozen) with medical treatment. Among the selected studies, there was a non-significant trend toward the resolution of diarrhea following a single fresh FMT infusion compared with frozen FMT or medical treatment (odds ratio, 2.45; 95% CI, 0.78 to 7.71; $p=0.12$, $I^2=69\%$), but different forms and routes of FMT administration were shown to be equally efficacious. In conclusion, FMT is a promising treatment modality for recurrent CDI. However, the authors found limited data for the variability of FMT dose usages, small trial populations and time frames to assess the success or failure of treatment.

In 2019, a third meta-analysis (Tariq et al) evaluated the efficacy of FMT as a treatment option for recurrent CDI on the basis of results from open-label studies and placebo controlled clinical trials. The authors wanted to investigate their observations on FMT cure rates for CDI being high in observational studies (e.g., >90%) but then appear to be consistently lower in open-label studies and clinical trials. Thirteen studies were included for evaluation, including six placebo-controlled RCTs and seven open-label studies. Out of 610 patients receiving FMT, 439 patients achieved clinical cure (76.1%; 95% confidence interval [CI]: 66.4% to 85.7%); study heterogeneity was significant ($I^2=91.35\%$). Cure rates were found to be lower in randomized trials (139/216, 67.7%; 95% CI: 54.2% to 81.3%) vs open-label studies (300/394, 82.7%; 95% CI: 71.1% to 94.3%; $p < 0.001$). Subgroup meta-analysis by FMT route of administration indicated lower cure rates with enema than colonoscopy (66.3% vs 87.4%; $p < 0.001$). However, no differences between colonoscopy and oral delivery were detected (87.4% to 81.4%; $p=0.17$). Lower cure rates were observed for studies that included both recurrent and refractory CDI than those that only included patients with recurrent CDI (63.9% vs 79%; $p < 0.001$). The authors concluded that colonoscopy and oral route are more effective than enema for stool delivery and the efficacy seems to be higher for recurrent than for refractory CDI.

A recent UpToDate (2020) review summarizes FMT as an effective treatment for recurrent (≥ 3 recurrences) CDI. The authors concluded that transplantation of stool microbiota from healthy individuals to patients with recurrent CDI can break the cycle of CDI recurrence. In addition, the initial results from the FMT National Registry published in Gastroenterology 2021 showed a 1 month cure rate

of over 90% in 200 patients who received just 1 FMT. At a 6 month follow-up, only 4% of patient had a recurrence of CDI and less than 1% had a complication of treatment includes IBS or newly diagnosed IBD. However, to minimize risk of infection, rigorous screening of potential healthy stool donors for occult pathogens must be done. The optimal approach for administration is uncertain. If feasible oral capsules should be tried first, then colonoscopy followed by enema retention and finally NJ or ND tube for patients who cannot undergo FMT via the alternate routes.

In 2016, the U.S. Food and Drug Administration (FDA) issued an updated draft guidance on enforcement policy regarding investigational new drug requirements for use of FMT to treat CDI not responsive to standard therapies. The draft guidance is similar to the 2013 guidance and states that the FDA is continuing to consider how to regulate FMT and that, during this interim period, the agency will use enforcement discretion regarding the use of fecal transplant to treat treatment-resistant CDI. The FDA requires that physicians obtain adequate informed consent from patients or their legal representative before performing the intervention. The document also stated that selective enforcement does not apply to the use of fecal transplant for treating conditions other than treatment-resistant CDI.

In 2019, the FDA issued a safety regarding the use of FMT due to the potential risk of serious adverse reactions or life-threatening infections caused by due to the transmission of multi-drug resistant organisms (MDROs). Two immunocompromised individuals received investigational FMT and developed invasive infections caused by the transmission of extended-spectrum beta-lactamase-producing *Escherichia coli*. One of the affected individuals died. The donor stool used in each patient's FMT procedures had not been tested for extended-spectrum beta-lactamase (ESBL)-producing gram-negative organisms prior to use. Follow-up testing verified donor stool was positive for MDROs identical to the organisms isolated from the two patients. Due to these events, the FDA has determined that the following additional protections are required for any investigational use of FMT:

- 1) Donor screening that specifically addresses risk factors for colonization with MDROs and exclusion of individuals at higher risk of colonization with MDROs such as health care workers, persons who have recently been hospitalized or discharged from long-term care facilities, persons who regularly attend outpatient medical or surgical clinics, and persons who have recently engaged in medical tourism.
- 2) MDRO testing of donor stool and exclusion of stool testing positive for MDROs. At the minimum, tests should include: extended-spectrum beta-lactamase-producing enterobacteriaceae, vancomycin-resistant enterococci, carbapenem-resistant enterobacteriaceae and methicillin-resistant *Staphylococcus aureus*.
- 3) All FMT products currently in storage for future use must be quarantined until donor MDRO carriage risk can be assessed and FMT products are tested and found negative for MDROs.
- 4) The informed consent process for FMT treatment subjects should describe the risk of MDRO transmission and infection and the measures being implemented for donor screening and stool testing.

Lastly, the American College of Gastroenterology (2021) published guidelines on diagnosis, treatment, and prevention of CDI. The guidelines addressed fecal microbiota transplant for treatment of three or more CDI recurrences, as follows: "For the treatment of one to two CDI recurrences, the guidelines recommend: tapering/pulsed-dose vancomycin for patients experiencing a first recurrence after an initial course of fidaxomicin, vancomycin, or metronidazole (strong recommendation, very low quality of evidence). In addition, Fidaxomicin for patients experiencing a first recurrence after an initial course of vancomycin or metronidazole. FMT be considered for patients with severe and fulminant CDI refractory to antibiotic therapy, in particular, when patients are deemed poor surgical candidates." In addition, fecal transplant is now recommended to be an important treatment consideration for refractory

antibiotic resistant CDI for patients who have underlying ulcerative colitis (UC) given the risk of exacerbation of the patient's underlying UC and the risk of complications from fulminant CDI in this particular patient population.

The Infectious Diseases Society of America and Society for Healthcare Epidemiology of America updated their clinical practice guidelines in 2018 for the diagnosis and treatment of CDI in children and adults. For pediatric patients fecal microbiota transplantation may be used after multiple recurrences of CDI following standard antibiotic treatments, this is considered a weak recommendation with very low quality of evidence. In adult patients, fecal microbiota transplantation is strongly recommended for patients with multiple recurrences of CDI who have failed appropriate antibiotic treatments as there is moderate quality of evidence.

No recommendations were found from the U.S. Preventive Services Task Force.

Applicable Coding

CPT Codes

- | | |
|--------------|--|
| 0780T | Instillation of fecal microbiota suspension via rectal enema into lower gastrointestinal tract |
| 44705 | Preparation of fecal microbiota for instillation, including assessment of donor specimen |

HCPCS Codes

- | | |
|--------------|---|
| G0455 | Preparation with instillation of fecal microbiota by any method, including assessment of donor specimen |
|--------------|---|

Not covered:

- | | |
|--------------|-------------------------------------|
| J1440 | Fecal microbiota, live - jslm, 1 ml |
|--------------|-------------------------------------|

ICD-10 Codes

- | | |
|---------------|--|
| A04.7 | Enterocolitis due to <i>Clostridium difficile</i> |
| A04.71 | Enterocolitis due to <i>Clostridium difficile</i> , recurrent |
| A04.72 | Enterocolitis due to <i>Clostridium difficile</i> , not specified as recurrent |

References:

1. American Journal of Gastroenterology (AJG). (April, 2013). "Guidelines for Diagnosis, Treatment, and Prevention of *Clostridium difficile* Infections" American Journal of Gastroenterology: April 2013 - Volume 108 - Issue 4 - p 478-498 doi: 10.1038/ajg.2013.4. Accessed September 18, 2020. Available at: https://journals.lww.com/ajg/Fulltext/2013/04000/Guidelines_for_Diagnosis,_Treatment,_and_Prevention_of_Clostridium_difficile_Infections.6.aspx#pdf-link
2. American Society of Colon and Rectal Surgeons (2015). Clinical Practice Guidelines. "Practice Parameters for the Management of *Clostridium difficile* Infection". Accessed September 18, 2020. Available at: https://fascrs.org/ascrs/media/files/downloads/Clinical%20Practice%20Guidelines/practice_parameter_clostridium_difficile_infection.pdf
3. Aroniadis, O.C., Brandt, L.J., Oneto, C., Feuerstadt, P., Sherman, A., Wolkoff, A.W., Kassam, Z., Sadovsky, R.G., Elliott, R.J., Budree, S. and Kim, M., 2019. Faecal microbiota transplantation for diarrhoea-predominant irritable bowel syndrome: a double-blind, randomised, placebo-controlled trial. The lancet Gastroenterology & hepatology, 4(9), pp.675-685.
4. Costello SP, Hughes PA, Waters O et al. Effect of Fecal Microbiota Transplantation on 8-Week Remission in Patients With Ulcerative Colitis: A Randomized Clinical Trial. JAMA, 2019 Jan 16;321(2). PMID 30644982
5. El-Salhy M, Hausken T, Hatlebakk JG. Current status of fecal microbiota transplantation for irritable bowel syndrome. Neurogastroenterology & Motility. 2021 Nov;33(11):e14157.

6. Feuerstadt P, Boules M, Stong L, Dahdal DN, Sacks NC, Lang K, Nelson WW. Clinical complications in patients with primary and recurrent *Clostridioides difficile* infection: A real-world data analysis. *SAGE Open Med.* 2021 Jan 14;9:2050312120986733. doi: 10.1177/2050312120986733. PMID: 33505698; PMCID: PMC7812403.
7. Food and Drug Administration (FDA). (2016) Guidance for Industry: Enforcement Policy Regarding Investigational New Drug Requirements for Use of Fecal Microbiota for Transplantation to Treat *Clostridium difficile* Infection Not Responsive to Standard Therapies. Accessed September 10, 2020. Available at: <https://www.fda.gov/regulatory-information/search-fda-guidance-documents/enforcement-policy-regarding-investigational-new-drug-requirements-use-fecal-microbiota-0>.
8. Food and Drug Administration (FDA). (2019) Fecal Microbiota Transplantation: Safety Communication - Risk of Serious Adverse Reactions Due to Transmission of Multi-Drug Resistant Organisms. Accessed September 10, 2020 Available at: <https://www.fda.gov/safety/medical-product-safety-information/fecal-microbiota-transplantation-safety-communication-risk-serious-adverse-reactions-due>.
9. Kelly CR, Fischer M, Allegretti JR, LaPlante K, Stewart DB, Limketkai BN, Stollman NH. ACG Clinical Guidelines: Prevention, Diagnosis, and Treatment of *Clostridioides difficile* Infections. *Official journal of the American College of Gastroenterology* | ACG. 2021 Jun 1;116(6):1124-47.
10. Kelly CR, Yen EF, Grinspan AM, Kahn SA, Atreja A, Lewis JD, Moore TA, Rubin DT, Kim AM, Serra S, Nersesova Y, Fredell L, Hunsicker D, McDonald D, Knight R, Allegretti JR, Pekow J, Absah I, Hsu R, Vincent J, Khanna S, Tangen L, Crawford CV, Mattar MC, Chen LA, Fischer M, Arsenescu RI, Feuerstadt P, Goldstein J, Kerman D, Ehrlich AC, Wu GD, Laine L. Fecal Microbiota Transplantation Is Highly Effective in Real-World Practice: Initial Results From the FMT National Registry. *Gastroenterology.* 2021 Jan;160(1):183-192.e3. doi:10.1053/j.gastro.2020.09.038. Epub 2020 Oct 1. PMID: 33011173; PMCID: PMC8034505.
11. Khan MY, Dirweesh A, Khurshid T et al. Comparing fecal microbiota transplantation to standard of-care treatment for recurrent *Clostridium difficile* infection: a systematic review and metaanalysis. *Eur J Gastroenterol Hepatol*, 2018 Aug 24;30(11). PMID 30138161
12. Khoruts A, Brandt LJ. Fecal microbiota transplant: a rose by any other name. *Official journal of the American College of Gastroenterology* | ACG. 2019 Jul 1;114(7):1176.
13. Luo Y, Lucas AL, Grinspan AM. Fecal transplants by colonoscopy and capsules are cost-effective strategies for treating recurrent *Clostridioides difficile* infection. *Digestive diseases and sciences.* 2020 Apr;65(4):1125-33.
14. Malnick, S.D., Fisher, D., Somin, M. and Neuman, M.G., 2021. Treating the metabolic syndrome by fecal transplantation—current status. *Biology*, 10(5), p.447.
15. McDonald LC, Gerding DN, Johnson S et al. Clinical Practice Guidelines for *Clostridium difficile* Infection in Adults and Children: 2017 Update by the Infectious Diseases Society of America (IDSA) and Society for Healthcare Epidemiology of America (SHEA). *Clin. Infect. Dis.*, 2018 Feb 21;66(7). PMID 29462280
16. Quraishi MN, Widlak M, Bhala N, et al. Systematic review with meta-analysis: the efficacy of faecal microbiota transplantation for the treatment of recurrent and refractory *Clostridium difficile* infection. *Aliment Pharmacol Ther.* Sep 2017;46(5):479-493. PMID 28707337
17. Ramai D, Zakhia K, Ofosu A, Ofori E, Reddy M. Fecal microbiota transplantation: donor relation, fresh or frozen, delivery methods, cost-effectiveness. *Annals of gastroenterology.* 2019 Jan;32(1):30.
18. Rosenbaum JT. Just another crappy commentary: the future of fecal microbiota transplantation. *Expert review of clinical immunology.* 2019 Oct 3;15(10):987-9.
19. Rossen NG, Fuentes S, van der Spek MJ, Tijssen JG, Hartman JH, Duflou A, Löwenberg M, van den Brink GR, Mathus-Vliegen EM, de Vos WM, Zoetendal EG. Findings from a randomized controlled trial of fecal transplantation for patients with ulcerative colitis. *Gastroenterology.* 2015 Jul 1;149(1):110-8.
20. Rubin DT, Ananthakrishnan AN, Siegel CA et al. ACG Clinical Guideline: Ulcerative Colitis in Adults. *Am. J. Gastroenterol.*, 2019 Mar 7;114(3). PMID 30840605 32. McDonald LC, Gerding DN, Johnson S et al. Clinical Practice Guidelines for *Clostridium difficile* Infection in Adults and Children: 2017 Update by the Infectious Diseases Society of America (IDSA) and Society for Healthcare Epidemiology of America (SHEA). *Clin. Infect. Dis.*, 2018 Feb 21;66(7). PMID 29462280
21. Santiago M, Eysenbach L, Allegretti J, Aroniadis O, Brandt LJ, Fischer M, Grinspan A, Kelly C, Morrow C, Rodriguez M, Osman M. Microbiome predictors of dysbiosis and VRE decolonization in patients with recurrent *C. difficile* infections in a multi-center retrospective study. *AIMS microbiology.* 2019;5(1):1.
22. Forogh B, Ghaseminejad Raeini A, Jebeli Fard R, Mirghaderi P, Nakhostin-Ansari A, Nakhostin-Ansari N, Bahari H, Hoveidaei AH. Efficacy of trigger point dry needling on pain and function of the hip joint: a systematic review of randomized clinical trials. *Acupuncture in Medicine.* 2024 Apr;42(2):63-75.
23. Stripling J, Rodriguez M. Current Evidence in Delivery and Therapeutic Uses of Fecal Microbiota Transplantation in Human Diseases—*Clostridium difficile* Disease and Beyond. *The American Journal of the Medical Sciences.* 2018 Nov 1;356(5):424-32.
24. Tariq R, Pardi DS, Bartlett MG et al. Low Cure Rates in Controlled Trials of Fecal Microbiota Transplantation for Recurrent *Clostridium difficile* Infection: A Systematic Review and Metaanalysis. *Clin. Infect. Dis.*, 2019 Apr 9;68(8). PMID 30957161

25. UpToDate® (2023) "Fecal microbiota transplantation for treatment of recurrent Clostridioides (formerly Clostridium) difficile infection". Topic 2604. Version 60.0. Topic last updated May 18, 2023. Literature current through June 2023. Accessed July 3, 2023. Available at: https://www.uptodate.com/contents/fecal-microbiota-transplantation-for-treatment-of-recurrent-clostridioides-formerly-clostridium-difficile-infection?search=fecal%20microbiota%20transplant&source=search_result&selectedTitle=1~37&usage_type=default&display_rank=1

Disclaimer:

This document is for informational purposes only and should not be relied on in the diagnosis and care of individual patients. Medical and Coding/Reimbursement policies do not constitute medical advice, plan preauthorization, certification, an explanation of benefits, or a contract. Members should consult with appropriate health care providers to obtain needed medical advice, care, and treatment. Benefits and eligibility are determined before medical guidelines and payment guidelines are applied. Benefits are determined by the member's individual benefit plan that is in effect at the time services are rendered.

The codes for treatments and procedures applicable to this policy are included for informational purposes. Inclusion or exclusion of a procedure, diagnosis or device code(s) does not constitute or imply member coverage or provider reimbursement policy. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage of these services as it applies to an individual member.

U of U Health Plans makes no representations and accepts no liability with respect to the content of any external information cited or relied upon in this policy. U of U Health Plans updates its Coverage Policies regularly, and reserves the right to amend these policies and give notice in accordance with State and Federal requirements.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from U of U Health Plans.

"University of Utah Health Plans" and its accompanying logo, and its accompanying marks are protected and registered trademarks of the provider of this Service and or University of Utah Health. Also, the content of this Service is proprietary and is protected by copyright. You may access the copyrighted content of this Service only for purposes set forth in these Conditions of Use.

© CPT Only – American Medical Association