# HEALTHY U MEDICAID

## MEDICAL PHARMACY PRIOR AUTHORIZATION REQUEST FORM TECARTUS®

For authorization, please answer each question and fax this form PLUS chart notes back to the Healthy U Prior Authorization Department at 801-213-1547.

## Failure to submit clinical documentation to support this request will result in a dismissal of the request.

If you have medical pharmacy prior authorization questions, please call for assistance: 833-981-0212

Disclaimer: Prior authorization request forms are subject to change in accordance with Federal and State notice requirements.

Date:	Member Name:	ID#:
DOB:	Gender:	Physician:
Office Phone:	Office Fax:	Office Contact:
Height/Weight:		HCPCS Code:

Member must try formulary preferred drugs before a request for a non-preferred drug may be considered. If treatment with preferred products has not been successful, you must submit which preferred products have been tried, dates of treatment, and reason for failure. Reasons for failure must meet the Health Plan medical necessity criteria.

**Product being requested:** 
□ Tecartus<sup>®</sup> (brexucabtagene autoleucel)

Dosing/Frequency:\_\_\_

Questions	Yes	No	Comments/Notes
MANTLE CELL LYMPHOM		NO	connients/Notes
1. Is the member 18 years of age or older?	1		
<ol> <li>Does the member have a diagnosis of relapse or refractory mantle</li> </ol>			Please provide
cell lymphoma?			documentation
3. Was the member refractory, or had no response, to all of the			Please provide
following:			documentation
<ul> <li>Anthracycline or bendamustine-containing regimen</li> </ul>			
<ul> <li>Anti-CD20 therapy, such as rituximab</li> </ul>			
<ul> <li>Bruton's Tyrosine Kinase (BTK) inhibitors, such as ibrutinib or acalabrutinib?</li> </ul>			
4. Does the member have a histological confirmation of cyclin D1			Please provide
overexpression or presence of the translocation t(11;14)?			documentation
5. Does the member have at least one measurable lesion per the			Please provide
Lugano Classification?			documentation
• Lymph nodes: the longest diameter in axial plane is >1.5 cm			
• Extranodal lesions: the longest diameter in axial plane is > 1.0			
cm			
6. Does the member have adequate bone marrow reserve with all of			Please provide
the following:			documentation
<ul> <li>Platelet count ≥ 75,000/µL</li> </ul>			
<ul> <li>Absolute neutrophil count (ANC) ≥ 1,000 cells/µL</li> </ul>			
<ul> <li>Absolute lymphocyte count (ALC) ≥ 100 cells/µL</li> </ul>			

7. Does the member have a Karnofski score ≥ 70 or Eastern		Please provide
Cooperative Oncology Group (ECOG) score < 2?		documentation
8. Does the member have a documented ability to tolerate a		Please provide documentation
lymphodepleting chemotherapy regimen of cyclophosphamide 500mg/m <sup>2</sup> intravenously and fludarabine 30mg/m <sup>2</sup> intravenously		documentation
on the fifth, fourth, and third days before Tecartus <sup>™</sup> infusion? 9. Does the member have adequate and stable renal, cardiac, and		Please provide
pulmonary function with all of the following:		documentation
<ul> <li>Creatinine clearance ≥ 60mL/min</li> </ul>		uocumentation
• Cardiac ejection fraction $\geq$ 50% and no evidence of pericardial		
effusion determined by an echocardiogram		
<ul> <li>Baseline oxygen saturation &gt; 92% on room air?</li> </ul>		
10. Dose the member have any of the following:		
<ul> <li>Previous allogeneic hematopoietic cell transplant (HSCT)</li> </ul>		
<ul> <li>Prior treatment with chimeric antigen receptor therapy or</li> </ul>		
other genetically modified T-cell therapy; or is being considered		
for treatment with any other gene therapy		
• Presence of fungal, bacterial, viral, or other infection that is		
uncontrolled requiring IV antimicrobials for management prior		
to Tecartus <sup>™</sup> infusion		
<ul> <li>Active inflammatory disorders</li> </ul>		
<ul> <li>Active hepatitis B (HBsAG positive) or hepatitis C (anti-HCV</li> </ul>		
positive) virus, if viral load is detectable		
<ul> <li>History of central nervous system lymphoma</li> </ul>		
<ul> <li>Active central nervous system (CNS) lymphoma or CNS</li> </ul>		
disorders by imaging		
<ul> <li>Detectable malignant cells in the cerebrospinal fluid or brain</li> </ul>		
metastases?		
11. Has the member received live vaccines within 6 weeks prior to the		
start of lymphodepleting chemotherapy?		
12. Will Tecartus <sup>™</sup> be given at a certified center to administer		
Tecartus™?	 	
13. Has the member and the requesting provider enrolled in the		
Yescarta <sup>®</sup> and Tecartus <sup>™</sup> REMS program?		
<b>B-CELL PRECURSOR ACUTE LYMPHOBLASTI</b> 1. Has the member received a previous treatment course of the		Please provide
requested medication or another CD19-directed chimeric antigen		documentation
receptor (CAR-T) therapy, or any prior CD19 directed therapy other		documentation
than blinatumomab?		
2. Does the member have Philadelphia chromosome-negative disease		Please provide
that is relapsed or refractory as defined as one of the following:		documentation
<ul> <li>Primary refractory disease; or</li> </ul>		
<ul> <li>First relapse with remission of 12 months or less; or</li> </ul>		
<ul> <li>Relapsed or refractory disease after at least 2 previous lines of</li> </ul>		
systemic therapy; <i>or</i>		
<ul> <li>Relapsed or refractory disease after allogeneic stem cell</li> </ul>		
transplant (allo-SCT)		
· · · · ·		

#### **Confidentiality Notice**

This document and any accompanying document contain confidential information and is intended for the use of the individual or entity named on this transmission sheet. If you are not the intended recipient, you are hereby notified that any disclosure, copying, distribution or the taking of any action in reliance on the contents of this information is strictly prohibited and the document should be returned to this office immediately. If you have received this facsimile in error, please notify us by telephone immediately and destroy document received.

<ul> <li>3. Does the member have Philadelphia chromosome-positive disease and meets either of the following: <ul> <li>Relapsed or refractory disease despite treatment with at least 2 different tyrosine kinase inhibitors (TKIs) (e.g., bosutinib, dasatinib, imatinib, nilotinib, ponatinib); or</li> <li>The member is intolerant to TKI therapy</li> </ul> </li> </ul>			Please provide documentation		
4. Does the member have morphological disease in the bone marrow?			Please provide documentation		
5. Does the member have an ECOG performance status of 0 to 2?			Please provide documentation		
6. Does the member have adequate and stable kidney, liver, pulmonary, and cardiac function?			Please provide documentation		
7. Does the member have active hepatitis B, active hepatitis C, or any active uncontrolled infection?			Please provide documentation		
8. Does the member have active graft versus host disease?					
9. Does the member have an active inflammatory disorder?					
What medications and/or treatment modalities have been tried in the past for this condition? Please document name of treatment, reason for failure, treatment dates, etc.					

Additional information:

Physician Signature:

\*\*Failure to submit clinical documentation to support this request will result in a dismissal of the request.\*\*

Policy: PHARM-HU-M031 Origination Date: 01/01/2022 Reviewed/Revised Date: 04/19/2023 Next Review Date: 04/19/2024 Current Effective Date: 05/01/2023

### **Confidentiality Notice**

This document and any accompanying document contain confidential information and is intended for the use of the individual or entity named on this transmission sheet. If you are not the intended recipient, you are hereby notified that any disclosure, copying, distribution or the taking of any action in reliance on the contents of this information is strictly prohibited and the document should be returned to this office immediately. If you have received this facsimile in error, please notify us by telephone immediately and destroy document received.