

HEALTHY U CHIP

PRIOR AUTHORIZATION REQUEST FORM

LEQVIO®

For authorization, please answer each question and fax this form PLUS chart notes back to the Healthy U CHIP Prior Authorization Department at 385-425-4052.

Failure to submit clinical documentation to support this request will result in a dismissal of the request.

If you have prior authorization questions, please call for assistance: 385-425-5094

Disclaimer: Prior Authorization request forms are subject to change in accordance with Federal and State notice requirements.

Date:	Member Name:	ID#:
DOB:	Gender:	Physician:
Office Phone:	Office Fax:	Office Contact:

Height/Weight:

Member must try formulary preferred drugs before a request for a non-preferred drug may be considered. If treatment with preferred products has not been successful, you must submit which preferred products have been tried, dates of treatment, and reason for failure. Reasons for failure must meet the Health Plan medical necessity criteria.

Product being requested: Leqvio® (inclisiran)

Dosing/Frequency: _____

If the request is for reauthorization, proceed to reauthorization section.

Questions	Yes	No	Comments/Notes
HETEROZYGOUS FAMILIAL HYPERCHOLESTEROLEMIA			
1. Is the request made by, or in consultation with, a cardiologist, endocrinologist, lipidologist, or a physician who focuses in the treatment of cardiovascular risk management and/or lipid disorders?	<input type="checkbox"/>	<input type="checkbox"/>	Please provide documentation
2. Does the member have a diagnosis of heterozygous familial hypercholesterolemia (HeFH) confirmed by either of the following: <ul style="list-style-type: none"> • Untreated LDL-C level \geq 190 mg/dL in adults • Untreated LDL-C level \geq 160 mg/dL and tendon xanthoma in members < 20 years of age 	<input type="checkbox"/>	<input type="checkbox"/>	Please provide documentation
3. Does the member have one of the following: <ul style="list-style-type: none"> • Genetic confirmation testing that demonstrates LDL-R mutation, LDLRAP1 mutation, familial defective apo B100, or a PCSK9 mutation • A diagnosis confirmed by the World Health Organization/Dutch Lipid Network Criteria resulting in a score > 8 points • A diagnosis meeting the threshold for definite or possible/probable familial hypercholesterolemia per Simon Broome Criteria • Arcus senilis if < 45 years of age 	<input type="checkbox"/>	<input type="checkbox"/>	Please provide documentation

<ul style="list-style-type: none"> • Have a first degree relative with similarly elevated LDL-C, early ASCVD (<55 years of age for men, <65 years of age for women), tendon xanthoma, or corneal arcus 			
4. Does clinical documentation show a recent baseline low-density lipoprotein cholesterol (LDL-C) level?	<input type="checkbox"/>	<input type="checkbox"/>	Please provide documentation
5. Has the member failed to reach target LDL-C when on high-intensity statin therapy or maximally tolerated statin therapy for at least 8 continuous weeks and LDL-C remains ≥ 100 mg/dL?	<input type="checkbox"/>	<input type="checkbox"/>	Please provide documentation
6. Does the member show LDL-C is unresponsive to standard PCSK9 therapy for an adequate duration (i.e., 3-6 months)? <ul style="list-style-type: none"> • Documentation must show trial and failure to Repatha® 	<input type="checkbox"/>	<input type="checkbox"/>	Please provide documentation
7. Will Leqvio® be used concurrently with a maximally tolerated statin therapy?	<input type="checkbox"/>	<input type="checkbox"/>	Please provide documentation
8. If the member has a contraindication to all statins, does documentation show one of the following: <ul style="list-style-type: none"> • Active liver disease • Diagnosis or history of rhabdomyolysis • Pregnant or nursing mothers • Allergic reaction with rash and/or anaphylactic symptoms 	<input type="checkbox"/>	<input type="checkbox"/>	Please provide documentation
9. If the member has a hypersensitivity to statins, does documentation show all of the following: <ul style="list-style-type: none"> • Inability to tolerate at least 2 different statins at the lowest starting dose • Intolerance associated with confirmed, intolerable statin-related adverse effects or significant biomarker abnormalities • Symptom and/or biomarker resolution upon discontinuation • Attestation that adverse effects are not attributable to drug-drug interactions or recognized conditions that can cause similar changes (e.g., hypothyroidism) • Intolerance persists despite trials of all the following: low dose of same or different statin, statin is dosed intermittently, and alternate cholesterol lowering medications such as ezetimibe or a bile-acid sequestrant such as colesevelam is used 	<input type="checkbox"/>	<input type="checkbox"/>	Please provide documentation
10. Is the member at least 80% compliant for at least 6 months with their baseline therapy (i.e., statins, ezetimibe)?	<input type="checkbox"/>	<input type="checkbox"/>	Please provide documentation
11. Has the provider addressed lifestyle modifications (i.e., a heart healthy diet, the importance of exercise, and smoking cessation) been completed?	<input type="checkbox"/>	<input type="checkbox"/>	Please provide documentation
12. Will the member be concurrently receiving any of the following medications in combination with Leqvio®: <ul style="list-style-type: none"> • Praluent® (alirocumab) • Repatha® (evolocumab) • Nexletol® (bempedoic acid) • Nexlizet® (bempedoic acid and ezetimibe) 	<input type="checkbox"/>	<input type="checkbox"/>	Please provide documentation
HIGH RISK OF ATHEROSCLEROTIC CARDIOVASCULAR DISEASE (ASCVD)			
1. Is the request made by, or in consultation with, a cardiologist, endocrinologist, lipidologist, or a physician who focuses in the treatment of cardiovascular risk management and/or lipid disorders?	<input type="checkbox"/>	<input type="checkbox"/>	Please provide documentation

<p>2. Does the member have a diagnosis of high risk atherosclerotic cardiovascular disease (ASCVD) with one of the following:</p> <ul style="list-style-type: none"> • History of myocardial infarction • Non-hemorrhagic stroke • Symptomatic peripheral artery disease • Acute coronary syndromes • Coronary artery disease • Stable or unstable angina • Coronary or other arterial revascularization • Transient ischemic attack • Diabetes • 10-year Framingham risk score of 20% or higher 	<input type="checkbox"/>	<input type="checkbox"/>	Please provide documentation
<p>3. Does clinical documentation show a recent baseline low-density lipoprotein cholesterol (LDL-C) level?</p>	<input type="checkbox"/>	<input type="checkbox"/>	Please provide documentation
<p>4. Has the member failed to reach target LDL-C when on high-intensity statin therapy or maximally tolerated statin therapy for at least 8 continuous weeks and LDL-C remains ≥ 70 mg/dL?</p>	<input type="checkbox"/>	<input type="checkbox"/>	Please provide documentation
<p>5. Does the member show LDL-C is unresponsive to standard PCSK9 therapy for an adequate duration (i.e., 3-6 months)?</p> <ul style="list-style-type: none"> • Documentation must show trial and failure to Repatha® 	<input type="checkbox"/>	<input type="checkbox"/>	Please provide documentation
<p>6. Will Leqvio® be used concurrently with a maximally tolerated statin therapy?</p>	<input type="checkbox"/>	<input type="checkbox"/>	Please provide documentation
<p>7. If the member has a contraindication to all statins, does documentation show one of the following:</p> <ul style="list-style-type: none"> • Active liver disease • Diagnosis or history of rhabdomyolysis • Pregnant or nursing mothers • Allergic reaction with rash and/or anaphylactic symptoms 	<input type="checkbox"/>	<input type="checkbox"/>	Please provide documentation
<p>8. If the member has a hypersensitivity to statins, does documentation show all of the following:</p> <ul style="list-style-type: none"> • Inability to tolerate at least 2 different statins at the lowest starting dose • Intolerance associated with confirmed, intolerable statin-related adverse effects or significant biomarker abnormalities • Symptom and/or biomarker resolution upon discontinuation • Attestation that adverse effects are not attributable to drug-drug interactions or recognized conditions that can cause similar changes (e.g., hypothyroidism) • Intolerance persists despite trials of all the following: low dose of same or different statin, statin is dosed intermittently, and alternate cholesterol lowering medications such as ezetimibe or a bile-acid sequestrant such as colesevelam is used 	<input type="checkbox"/>	<input type="checkbox"/>	Please provide documentation
<p>9. Is the member at least 80% compliant for at least 6 months with their baseline therapy (i.e., statins, ezetimibe)?</p>	<input type="checkbox"/>	<input type="checkbox"/>	Please provide documentation
<p>10. Has the provider addressed lifestyle modifications (i.e., a heart healthy diet, the importance of exercise, and smoking cessation) been completed?</p>	<input type="checkbox"/>	<input type="checkbox"/>	Please provide documentation

11. Will the member be concurrently receiving any of the following medications in combination with Leqvio®: <ul style="list-style-type: none"> • Praluent® (alirocumab) • Repatha® (evolocumab) • Nexletol® (bempedoic acid), • Nexlizet® (bempedoic acid and ezetimibe) 	<input type="checkbox"/>	<input type="checkbox"/>	Please provide documentation
REAUTHORIZATION			
1. Is the request for reauthorization of therapy?	<input type="checkbox"/>	<input type="checkbox"/>	
2. Does documentation indicate an adequate reduction in LDL-C defined by one of the following: <ul style="list-style-type: none"> • ≥ 40% reduction in LDL-C level compared to baseline or reduction to LDL goal in members with a diagnosis of ASCVD • Reduction in LDL-C level compared to baseline in members with a diagnosis of HeFH 	<input type="checkbox"/>	<input type="checkbox"/>	Please provide documentation
3. Is member adherent to concurrent statin therapy at the maximum tolerated dose?	<input type="checkbox"/>	<input type="checkbox"/>	Please provide documentation
4. Is member adherent to lifestyle modifications (i.e., a heart healthy diet, the importance of exercise, and smoking cessation)?	<input type="checkbox"/>	<input type="checkbox"/>	Please provide documentation
What medications and/or treatment modalities have been tried in the past for this condition? Please document name of treatment, reason for failure, treatment dates, etc.			
Additional information:			
Physician Signature:			

**** Failure to submit clinical documentation to support this request will result in a dismissal of the request. ****

Policy: PHARM-CHIP-130
 Origination Date: 07/01/2024
 Reviewed/Revised Date:
 Next Review Date:
 Current Effective Date: 07/01/2024

Confidentiality Notice

This document and any accompanying document contain confidential information and is intended for the use of the individual or entity named on this transmission sheet. If you are not the intended recipient, you are hereby notified that any disclosure, copying, distribution or the taking of any action in reliance on the contents of this information is strictly prohibited and the document should be returned to this office immediately. If you have received this facsimile in error, please notify us by telephone immediately and destroy document received.