

PRIOR AUTHORIZATION REQUEST FORM

OSTEOPOROSIS MEDICATIONS

Evenity®, teriparatide, Tymlos®

For authorization, please answer each question and fax this form PLUS chart notes back to the U of U Health Plans Prior Authorization Department at 888-509-8142.

Failure to submit clinical documentation to support this request will result in a dismissal of the request.

If you have prior authorization questions, please call for assistance: University of Utah Health Employees: 855-856-5690, Individual & Family Plans: 855-869-4769. Commercial Groups: 855-859-4892. MHC 855-885-7695

	individual & Family Flans : 655-665 4705, Commercial Groups. 655-655-4652, Wire 655-665-7655						
Dis	claimer: Prior authorization request for	orms are subject to change in accord	lance wi	th Fede	ral and State notice requirements.		
		T					
Dat	te:	Member Name:		ID#:			
DO	B:	Gender:		Physi	cian:		
Office Phone:		Office Fax:	Office		e Contact:		
Hei	Height/Weight:						
Member must try formulary preferred drugs before a request for a non-preferred drug may be considered. If treatment with preferred products has not been successful, you must submit which preferred products have been tried, dates of treatment, and reason for failure. Reasons for failure must meet the Health Plan medical necessity criteria. Preferred: Evenity® (romosozumab), teriparatide, Tymlos® (abaloparatide) Dosing/Frequency:							
If the request is for reauthorization, proceed to reauthorization section							
	Questio	ns	Yes	No	Comments/Notes		
1.	Does the member have a docume	nted diagnosis of one of the			Please provide documentation		
	 following: Postmenopausal female with o Male with primary or hypogon Osteoporosis likely caused by stherapy? 	adal osteoporosis,			F		
2.	Postmenopausal female with oMale with primary or hypogon	adal osteoporosis, systemic glucocorticoid sk for fracture, defined by ure or bone mineral density orosis (T-score < -2.5), nd/or glucocorticoid use for at			Please provide documentation		
2.	 Postmenopausal female with of Male with primary or hypogon Osteoporosis likely caused by stherapy? Is the member considered high rismeeting one of the following: History of recent fragility fract measurement showing osteop History of previous fractures a least 3 months and osteopenia 2.5)? 	adal osteoporosis, systemic glucocorticoid sk for fracture, defined by ure or bone mineral density orosis (T-score ≤ -2.5), nd/or glucocorticoid use for at a (T-Score between -1 and - teoporosis, defined as one of gility fracture for fragility fracture based on					

alendronate, ibandronate, risedronate, intravenous zoledronic						
acid), unless contraindicated?						
 IV therapy (zoledronic acid) is required if the member is 						
unable to tolerate oral bisphosphonate or has an absorption						
disorder.						
5. Does documentation show a 24-month trial and failure of			Please provide documentation			
Prolia®, unless contraindicated?						
What medications and/or treatment modalities have been tried in the past for this condition? Please document						
name of treatment, reason for failure, treatment dates, etc.						
Additional information:						
Physician's Signature:						
,						

** Failure to submit clinical documentation to support this request will result in a dismissal of the request.**

Policy: PHARM- 054

Origination Date: 08/07/2017 Reviewed/Revised Date: 03/15/2023 Next Review Date: 03/15/2024 Current Effective Date: 04/01/2023

Confidentiality Notice